An Introduction to:

The Period of PURPLE Crying® a program of the National Center on Shaken Baby Syndrome

Veronica Wiesz vwiesz@pcand.org











a program of the Shaken Baby Syndrome

Magnitude of the Problem

- AHT is the leading cause of death from physical child abuse.
- AHT has an estimated incidence of 32-38 cases per 100,000 children per year in the first year of life, with fatality occurring in 25% of cases.
- Usually occurs in children less than one year old, peaking at 2-3 months of age.







Seconds of Frustration: Lifelong Cost

#1 trigger for AHT is inconsolable crying. This occurs as a reaction of frustration by the caregiver.


Consequences of Shaking

- Death
- Blindness
- Learning disabilities
- Physical disabilities
- Hearing loss
- Speech disabilities
- Cerebral Palsy
- Chronic seizures
- Behavior disorders





The Crying Curve



One of the BIGGEST STRESSORS for **PARENTS** is the normal period of increased crying all babies experience in the first few months of life.



What is PURPLE?



- Evidence-based program
- ► Goal 1: Support parents and caregiver's understanding of early increased crying
- ► Goal 2: Reduce the incidences of SBS/AHT
- Materials are educational, relevant, clear and memorable

The Letters in **PURPLE** Stand for



Your baby may cry more each week, the most in month 2, then less in months 3-5

Crying can come and go and you don't know why

Your baby may not stop crying no matter what you try

A crying baby may look like they are in pain, even when they are not


Crying can last as much as 5 hours a day, or more

Your baby may cry more in the late afternoon and evening

The word **Period** means that the crying has a beginning and an end.



Our History



Three Dose Exposure Approach



Targets parents and caregivers of newborns 0-2 weeks old.

 $\bullet \bullet \bullet$

Trained providers deliver education and materials to parents.

. . .

Hospital discharge setting,
Perinatal/Postnatal Classes, Home Visiting



Targets parents and caregivers of newborns 2 weeks to 3 months old.

• • •

Trained providers reinforce key program messages and deliver materials if necessary.

• • •

Home Visiting, Pediatric Well Baby Visits, Public & State Department of Health Programs



Targets the general population. Not set to timing of baby's birth.

• •

Dependent upon organization's capacity to educate the community.

• • •

"Normal" Ad Campaign, PURPLE Toolkit, CLICK for Babies, etc.

Supplementary Materials

New Dad's Handout

- Accompanies the *Period of PURPLE Crying* App+Booklet package, to support fathers as they care for their new baby.
- Discusses how dads can form loving attachments with their baby even when the crying can be frustrating and directs fathers to sections within the app that provide additional resources about soothing and coping.



New App Package Posters

- Highlights all the exciting resources and features of the app.
- Each partner will receive 3 complimentary posters with their next order and will have the ability to purchase more.
- Walks families through the app download process.



Please note: this resource is only available to organizations that are currently delivering the *Period of PURPLE Crying* program with fidelity.



Why the Period of PURPLE Crying?


Most
WIDELY
RESEARCHED
SBS/AHT program
in the
WORLD

Delivered to more than 3 MILLION FAMILIES by more than 2,000 HOSPITAL and public health systems worldwide

The program saved hospitals 18% in costs by reducing crying-related ER visits by 29%



Costs of AHT & PURPLE program



Medical Costs

A 2020 study revealed that the cost of medical care alone for a severely injured victim (2.5 mo. to age 17) was \$261,499. During that time, a total of \$106,607 was paid out-of-pocket by the family (Steinbeigle et al., 2020).

2

Projected Lifetime Costs

The total cost of AHT per mildly affected case is **over one million dollars**. From a societal perspective, the discounted average lifetime cost of a surviving AHT victim cost \$2.6 million and AHT death \$5.7 million (Miller et al., 2017).

(3)

Net Out-of-Pocket Saving

If existing prevention programs reduced AHT by **2%** they would yield a net out-of-pocket saving. *PURPLE* prevents **35%** of AHT events (Barr et al., 2018).

Cost-Effectiveness

Every \$1 spent on the *PURPLE* program had an associated **\$54** savings by society. This can be compared to Zero Tolerance (\$25/per \$1) & Child Safety Seat Laws (\$42/per \$1) (Beaulieu et al., 2019).



Thank You!! Questions? Veronica Wiesz vwiesz@pcand.org




Email



Training - Don't Shake









References:

- Barr, R. G., Barr, M., Rajabalic, F., Humphreys, C., Pike, I., Brante, R., . . . Singhalh, A. (2018). Eight-year outcome of implementation of abusive head trauma prevention. *Child Abuse & Neglect*, 106-114.
- Barr, R. G., Barr, M., Fujiwara, T., Conway, J., Catherine, N., & Brant, R. (2009). Do educational materials change knowledge and behaviour about crying and shaken baby syndrome? A randomized controlled trial. *Canadian Medical Association Journal*, 180(7), 727-733.
- Barr, R. G., Rivara, F. P., Barr, M., Cummings, P., Taylor, J., Lengua, L. J., & Meredith-Benitz, E. (2009). Effectiveness of Educational Materials Designed to Change Knowledge and Behaviors Regarding Crying and Shaken-Baby Syndrome in Mothers of Newborns: A Randomized, Controlled Trial. *Pediatrics*, 123(3), 972-980.
- Beaulieu, E., Rajabali, F., Zheng, A., & Pike, I. (2019). The lifetime costs of pediatric abusive head trauma and a cost-effectiveness analysis of the Period of Purple crying program in British Columbia, Canada. *Child Abuse & Neglect*.
- Berger, R. P., Fromkin, J. B., Stutz, H., Makoroff, K., Scribano, P. V., Feldman, K., . . . Fabio, A. (2011, October). *Abusive Head Trauma During a Time of Increased Unemployment: A Multicenter Analysis*. Retrieved from Pediatrics: https://pediatrics.aappublications.org/content/128/4/637
- Brazelton, T. B. (1990). Crying and colic. Infant Mental Health Journal.
- CDC. (2019, June 24). Deaths: Final Data for 2017. Retrieved from CDC: https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68_09-508.pdf
- CDC. (2020). A Journalist's Guide to Shaken Baby Syndrome. Retrieved from CDC: https://www.cdc.gov/violenceprevention/pdf/SBSMediaGuide.pdf
- Choudhary, A. K., Servaes, S., Slovis, T. L., Palusci, V. J., Hedlund, G. L., Narang, S. K., . . . Raissaki, M. (2018). Consensus statement on abusive head trauma in infants and young children. *Pediatric Radiology*.
- Gubbels, J., van der Put, C. E., Stams, G.-J. J., Prinzie, P. J., & Assink, M. (2021, April). Components Associated with the Effect of Home Visiting Programs on Child Maltreatment: A Meta-Analytic Review. Child Abuse & Neglect, 114.
- Joyce, T., & Huecker, M. R. (2019). Pediatric Abusive Head Truama (Shaken Baby Syndrome). StatPearls Publishing LLC.
- Lee, C., Barr, R., Catherine, N., & Wicks, A. (2007). Age-related incidence of publicly reported shaken baby syndrome cases: is crying a trigger for shaking? *Journal of developmental and behavioral pediatrics*, 288-93.
- McJunkin, J., & Phillips, J. (2020, April 9). McJunkin, Phillips: Young parents, babies need our support (Opinion). Retrieved from Charleston Gazette-Mail:

 https://www.wvgazettemail.com/opinion/op_ed_commentaries/mcjunkin-phillips-youngparents-babies-need-our-support-opinion/article_ee1612c5-3294-5222-b1e2-c4f0ce13a78e.html?utm_medium=social&utm_source=email&utm_campaign=user-share

References:

Miller, T. R., Steinbeigle, R., Lawrence, B. A., Peterson, C., Florence, C., Barr, M., & Barr, R. G. (2017). Lifetime Cost of Abusive Head Trauma at Ages 0-4, USA. Prevention Science.

Miller, T. R., Steinbeigle, R., Wicks, A., Lawrence, B. A., Barr, M., & Barr, R. G. (2014). Disability-Adjusted Life Year Burden of Abusive Head Trauma at Ages 0-4. *Pediatrics*.

NCSBS. (2019). Learn More. Retrieved from National Center on Shaken Baby Syndrome: https://www.dontshake.org/learn-more

Niederkrotenthaler, T., Xu, L., Parks, S. E., & Sugerman, D. E. (2013). Descriptive factors of abusive head trauma in young children—United States, 2000–2009. Child Abuse & Neglect, 446-455.

Reece, R. M., Dias, M. S., Barr, M., Russell, B. S., Barr, R. G., & Runyan, D. K. (2010, March). White Paper. Retrieved from Purplecrying.info: http://purplecrying.info/public_files/white_paper_SBSAHT_evidence_based_prevention.pdf

Reese, L. S., Heiden, E. O., Kim, K. Q., & Yang, J. (2014). Evaluation of Purple Crying, an Abusive Head Trauma Prevention Program. JOGNN, 752-761.

Runyan DK, et al. (2010) International variations in harsh child discipline. Unpublished Manuscript.

Schenck-Fontaine, A., & Gassman-Pines, A. (2020, May). Income inequality and child maltreatment risk during economic recession. Retrieved from ScienceDirect:

https://www.sciencedirect.com/science/article/pii/S019074091931014X

Steinbeigle, R., Barr, M., Barr, R. G., & Miller, T. R. (2020, May). Financial Impact of Abusive Head Trauma to Age 17. Child Abuse Review, 29(3), 208-17.

Zheng, L. (2020, March 21). Fort Worth Hospital Sees Spike in Severe Child Abuse Cases Over Last Week. Retrieved from NBCDFW: https://www.nbcdfw.com/news/local/fort-worth-hospital-sees-spike-in-severe-child-abuse-cases-over-last week/2336014/?fbclid=IwAR3Ve5Try0TY47DQ2pb tDrHYmf0IrrPQO7rhF8B6mMNCsKTGyPJD__KJ3w